



Airway Pressure Release Ventilation in Adult Patients With Acute Hypoxemic Respiratory Failure: A Systematic Review and Meta-Analysis*

Jolene Lim, MBBS (Hon), MSc (Dist)¹; Edward Litton, MBChB, FCICM, MSc, PhD^{1,2}

Objectives: To evaluate the efficacy and safety of airway pressure release ventilation in critically ill adults with acute hypoxemic respiratory failure.

Data Sources: A systematic literature search of MEDLINE via PUBMED, EMBASE, the Cochrane Library, published conference proceedings and abstracts, reference lists of eligible studies and review articles, and hand searches of relevant journals and trial registers.

Study Selection: Eligible studies included randomized controlled trials published between years 2000 and 2018, comparing airway pressure release ventilation to any ventilation mode, in critically ill adults with acute hypoxemic respiratory failure and reporting at least one mortality outcome.

Data Extraction: Screened citations were reviewed and extracted independently by two investigators onto a prespecified proforma.

Data Synthesis: There were 412 patients from seven randomized controlled trials included in the qualitative and quantitative data synthesis. Airway pressure release ventilation was associated with a significant mortality benefit (relative risk, 0.67; 95% CI, 0.48–0.94; $I^2 < 0.1\%$; $p = 0.97$) and improvement in day 3 P_{aO_2}/F_{iO_2} ratio (weighted mean difference, 60.4; 95% CI, 10.3–110.5). There was no significant difference in requirement to initiate rescue treatments including inhaled pulmonary vasodilators, prone positioning, or extracorporeal membrane oxygenation (relative risk, 0.51; 95% CI, 0.22–1.21; $I^2 = 64.7\%$; $p = 0.04$). The risk of barotrauma was only reported in three studies and did not

differ between groups (relative risk, 0.39; 95% CI, 0.12–1.19; $I^2 < 0.1\%$; $p = 0.99$).

Conclusions: In adult patients requiring mechanical ventilation for acute hypoxic respiratory failure, airway pressure release ventilation is associated with a mortality benefit and improved oxygenation when compared with conventional ventilation strategies. Given the limited number of patients enrolled in the available studies, larger multicenter studies are required to validate these findings. (*Crit Care Med* 2019; 47:1794–1799)

Key Words: acute hypoxemic respiratory failure; acute respiratory distress syndrome; airway pressure release ventilation

Mortality associated with acute respiratory failure requiring mechanical ventilation remains high, with an occurrence of approximately 50% in critically ill patients with hypoxemic respiratory failure (1). A pivotal randomized controlled trial (RCT) demonstrated improved outcomes with low tidal volume (LTV) ventilation but is now nearly 20 years old (2). Most subsequent studies examining alternative ventilatory strategies have failed to demonstrate benefit and there remains an unmet need for effective interventions in severely hypoxic patients (3–6).

Airway pressure release ventilation (APRV) is a pressure-controlled mode of ventilation that allows continuous, unrestricted spontaneous respiration in addition to intermittent mandatory ventilation (7, 8). As conventionally described, the majority of the respiratory cycle occurs at the higher of two set pressures (P_{high}), with brief periods of lower pressure (P_{low}), often set as zero. Theoretical benefits of APRV include increased lung recruitment, reduced atelectrauma, improved ventilation/perfusion (V/Q) matching and lower sedation and neuromuscular blockade requirements and hemodynamic benefits (7, 9). Observational studies have demonstrated an association with improved oxygenation and lower requirement for the initiation of rescue therapies including extracorporeal membrane oxygenation (ECMO) (10).

Recent RCTs suggest that APRV may improve survival (11, 12). However, concerns remain about the safety of APRV,

*See also p. 1817.

¹Department of Intensive Care, Fiona Stanley Hospital, Perth, WA, Australia.

²School of Medicine, University of Western Australia, Perth, WA, Australia.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website (<http://journals.lww.com/ccmjournal>).

The authors have disclosed that they do not have any potential conflicts of interest.

For information regarding this article, E-mail: ed_litton@hotmail.com

Copyright © 2019 by the Society of Critical Care Medicine and Wolters Kluwer Health, Inc. All Rights Reserved.

DOI: 10.1097/CCM.0000000000003972

especially the risk of volutrauma and barotrauma (13, 14). To address this, we performed a systematic review and meta-analysis with the aim of evaluating whether APRV, compared with other modes of mechanical ventilation, decreased all-cause mortality in critically ill adults with acute hypoxemic respiratory failure.

MATERIALS AND METHODS

The systematic review was prospectively registered on the International prospective register of systematic reviews (PROSPERO) (CRD42018114916) (15).

Eligibility Criteria

We included RCTs of adult patients admitted to ICU with acute hypoxemic respiratory failure requiring mechanical ventilation, in which APRV was compared with an alternate mode of ventilation. Observational studies, crossover studies, and studies that did not report mortality were excluded. Studies published before the year 2000 were also excluded as this was before the publication of the landmark Acute Respiratory Distress Syndrome Network trial which established LTV ventilation strategies as the current gold standard strategy and thus reduced inter-study

heterogeneity with regard to the control mode used as well as excluded modes of ventilation proven to be harmful (2).

Data Sources and Search Strategy

We performed a comprehensive search of MEDLINE via PUBMED, EMBASE, and the Cochrane Library from January 2000 to November 2018 without any language restrictions. Our search also included published conference proceedings and abstracts, reference lists of eligible studies and review articles, and hand searches of relevant journals and trial registers. The search was repeated prior to final analyses. The following key search terms (and their synonyms) were used as follows: APRV, hypoxemic respiratory failure, acute respiratory distress syndrome (ARDS), and critically ill.

Study Selection

Two reviewers (J.L., E.L.) independently conducted the primary search and screened the titles and abstracts of articles. The full text of potentially eligible studies were retrieved and independently assessed by both reviewers. The corresponding authors of relevant studies were contacted for further information if there was uncertainty over the eligibility of the study (16, 17). There were no major disagreements related to study eligibility criteria

or outcomes. Minor disagreements between the authors were resolved through discussions without the requirement for a mediator.

Data Extraction and Quality Assessment

A prespecified proforma was used to extract relevant data on the study design, patient demographics, intervention, control, and outcomes. Missing data were requested from the corresponding authors of the relevant studies. The risk of bias in individual studies was assessed using the Cochrane Risk of Bias tool (18).

Data Analyses

The primary outcome measure was the all-cause mortality at the longest reported time point. Secondary outcomes included risk of all-cause barotrauma defined as new pneumothorax, pneumomediastinum, or subcutaneous emphysema occurring after initiation of APRV or the comparator ventilation mode, occurrence of cardiac arrest, $\text{PaO}_2/\text{FiO}_2$ on day 3,




Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow diagram demonstrating study selection for inclusion in the quantitative meta-analysis. RCT = randomized controlled trial.

TABLE 1. Risk of Bias of Individual Studies

| References | Random Sequence Generation (Selection Bias) | Allocation Concealment (Selection Bias) | Blinding of Participants and Personnel (Performance Bias) | Blinding of Outcome Assessment (Detection Bias) | Incomplete Outcome Data (Attrition Bias) | Selective Reporting (Reporting Bias) | Other Bias |
|----------------------|---|---|---|---|--|--------------------------------------|--------------|
| Hirshberg et al (12) | Low risk | Low risk | High risk | Unclear risk | Low risk | Low risk | Low risk |
| Li et al (21) | Low risk | Unclear risk | High risk | Unclear risk | Low risk | Unclear risk | Low risk |
| Maxwell et al (22) | Low risk | Unclear risk | High risk | Unclear risk | Low risk | Low risk | Low risk |
| Putensen et al (23) | Unclear risk | Unclear risk | High risk | Unclear risk | Low risk | Low risk | Unclear risk |
| Varpula et al (24) | Low risk | Low risk | High risk | Unclear risk | Low risk | Low risk | Low risk |
| Zhou et al (11) | Low risk | Low risk | High risk | Unclear risk | Low risk | Low risk | Low risk |
| Patel et al (17) | Unclear risk | Unclear risk | High risk | Unclear risk | Unclear risk | Unclear risk | Unclear risk |

number of ventilator-free days to day 28, and requirement for rescue interventions defined as ECMO, proning, neuromuscular paralysis, recruitment maneuvers, high-frequency oscillatory ventilation, inhaled nitric oxide, and/or inhaled epoprostenol.

Where there were three or more studies with an outcome of interest, a quantitative synthesis via meta-analysis was conducted. Standardized mean differences (SMDs) and weighted mean differences (WMDs) for continuous outcomes and risk ratios (RRs) for binary outcomes were calculated with 95% CIs and two-sided *p* values for each outcome. SD were derived from SEMs or 95% CIs based on formulae from the Cochrane Handbook if not directly reported (19). The results were pooled and analyzed primarily using the random-effects model due to anticipated significant inter-study heterogeneity. Fixed-effects model analysis were also conducted and reported if results were substantially different. Heterogeneity was assessed using the chi-square test and *I*² statistic. Sensitivity analyses excluding studies assessed to have a high risk of bias were performed.

We prespecified subgroup analyses comparing APRV with studies using LTV as the comparator mode versus alternative ventilation modes, and studies that exclusively enrolled patients with ARDS versus more heterogeneous pathologies. Risk of publication bias was assessed using Egger's test and a funnel plot. Trial sequential analysis (TSA) was also conducted to assess the risk of false inferences assuming a relative risk reduction in mortality of 33% from a baseline rate of 30%, double-sided $\alpha = 5\%$, $\beta = 20\%$ (power of 80%) and heterogeneity based on model variance (20). Stata 14 (StataCorp, College Station, TX) was used for all analysis except TSA which was conducted using TSA software (Copenhagen Trial Unit, Center for Clinical Intervention Research, Rigshospitalet, Copenhagen, Denmark).

RESULTS

There were 162 citations initially identified for review and seven studies with a total of 412 participants (11, 12, 17, 21–24) were included in the qualitative and quantitative data analysis (Fig. 1).

The causes of acute hypoxemic respiratory failure included both pulmonary and extrapulmonary pathology, with the majority related to pneumonia or trauma. The mean Acute

Physiology and Chronic Health Evaluation II score was 20.26 (SD 5.4), and 308 patients (74.8%) were diagnosed with ARDS or acute lung injury (this was used in studies predating the Berlin definition [25]). The study and participant characteristics are provided in **Supplemental Table 1** (Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>). The overall quality of the studies was moderate (**Table 1**).

OUTCOMES

Mortality

APRV was associated with a significant decrease in all-cause mortality (RR, 0.67; 95% CI, 0.48–0.94) without significant heterogeneity ($I^2 < 0.1\%$; $p = 0.97$) (Fig. 2). There was no evidence of publication bias based on the funnel plot (Supplemental Fig. 1, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>) or Egger's test (0.49; 95% CI, –0.17 to 1.15; $p = 0.11$).

The results were similar when analysis excluded four studies at high risk of bias (RR, 0.61; 95% CI, 0.41–0.91; $I^2 < 0.1\%$; $p = 0.87$) (Supplemental Fig. 2, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>).

On subgroup analysis, the point estimate for the effect of APRV on mortality was similar when including studies exclusively with LTV as a comparator ventilation strategy (RR, 0.67; 95% CI, 0.47–0.96; $I^2 < 0.1\%$; $p = 0.94$) compared with the single study in which APRV was compared with an alternative ventilation strategy (pressure-controlled synchronized intermittent mandatory ventilation aiming for tidal volume 8–10 mL/kg), (RR, 0.67; 95% CI, 0.24–1.86) (Supplemental Fig. 3, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>) (24).

The point estimate for the effect of APRV on mortality was also similar for studies exclusively enrolling patients with ARDS versus more heterogeneous pathologies (RR, 0.70; 95% CI, 0.47–1.03; $I^2 < 0.1\%$; $p = 0.915$) versus (RR, 0.61; 95% CI, 0.31–1.17; $I^2 < 0.1\%$; $p = 0.74$), respectively (Supplemental Figs. 4 and 5, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>).

The results were also similar on sensitivity analysis, limited to studies reporting short-term mortality (longest available of 28-d, 30-d, ICU, and in-hospital mortality), (RR, 0.69; 95% CI,




Figure 2. Forest plot of studies comparing airway pressure release ventilation (APRV) versus alternative modes of mechanical ventilation on mortality. RR = relative risk.

0.49–0.98; $I^2 < 0.1\%$, $p = 0.91$) (Supplemental Fig. 6, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>).

TSA demonstrated that although the cumulative Z-curve surpassed the conventional 95% confidence boundary favoring APRV, the trial sequential monitoring boundaries adjusting for sequential testing were not crossed and the required information size of 602 participants was also not achieved (Supplemental Fig. 7, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>).

Oxygenation Rescue Strategies and Ventilation

Day 3 P_{aO_2}/F_{iO_2} was reported by five studies and was significantly higher in patients receiving APRV although there was




Figure 3. Forest plot of studies comparing airway pressure release ventilation (APRV) versus alternative modes of mechanical ventilation on P_{aO_2}/F_{iO_2} ratio on day 3. WMD = weighted mean difference.

significant inter-study heterogeneity (WMD, 60.4; 95% CI, 10.3–110.5; $I^2 = 93.4\%$; $p < 0.01$) (Fig. 3). The results were similar when assessing SMD in day 3 P_{aO_2}/F_{iO_2} (SMD, 0.82; 95% CI, 0.60–1.05; $I^2 = 87.8\%$; $p < 0.01$) (Supplemental Fig. 8, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>). There was no significant difference in requirement to initiate rescue treatments which was reported by four studies (RR, 0.51; 95% CI, 0.22–1.21; $I^2 = 64.7\%$; $p = 0.04$) (Supplemental Fig. 9, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>).

Safety

In total, three studies reported barotrauma with a total of 12 incidents, all of which were pneumothoraces (three in APRV group, nine in the control group) (11, 12, 22). The risk of barotrauma did not differ significantly between groups (RR, 0.39; 95% CI, 0.12–1.19; $I^2 < 0.1\%$; $p = 0.99$) (Supplemental Fig. 10, Supplemental Digital Content 1, <http://links.lww.com/CCM/E886>). None of the included studies reported the occurrence of cardiac arrest, volutrauma, or other serious adverse events.

DISCUSSION

In our systematic review of adult patients requiring mechanical ventilation for acute hypoxemic respiratory failure, APRV compared with other ventilator modalities was associated with lower mortality (RR, 0.67; 95% CI, 0.48–0.94). The findings were consistent when limiting the analysis to higher-quality studies, studies exclusively enrolling patients with ARDS, and studies exclusively comparing APRV to LTV ventilation strategies. APRV was also associated with a significant improvement in day 3 P_{aO_2}/F_{iO_2} ratio (WMD, 60.4; 95% CI, 10.3–110.5). The

number of barotrauma events was low and did not differ between treatment groups.

Several ventilation strategies have been used with the aim of improving oxygenation through the recruitment of atelectatic lung. However, some have been demonstrated to cause harm (3, 6). In our analysis, day 3 oxygenation was improved among patients ventilated with APRV, but a signal of harm was not detected. Rather, mortality appears to be lower with APRV, suggesting that while APRV also increases lung recruitment and oxygenation (8, 13), it potentially does so without increasing the risk of barotrauma or other deleterious consequences.

In our analysis, patients receiving APRV had higher mean airway pressure despite receiving lower peak airway pressures than the comparator group. Although these measures are surrogates for the risk associated with high transpulmonary pressure and assume peak pressures correlate with plateau pressures, these results suggest that the lower peak pressures and higher mean airway pressures achieved with APRV may reduce the risk of secondary lung injury while still allowing progressively greater recruitment. Mean measured tidal volumes in the APRV group (7.47 mL/kg [SD 1.22]) were also very similar to the comparator group (7.45 mL/kg [SD 1.27]) where all but one trial used LTV strategies (although only three studies reported expired tidal volume among the APRV patients).

Other mechanisms may contribute to a decreased risk of death with APRV. APRV may result in greater ventilator synchrony when compared with LTV and thus lower sedation requirement (7). Excess sedation in the comparator groups in APRV RCTs may contribute to increased mortality risk. The risk of requiring rescue therapy was also lower in patients receiving APRV, although statistical significance was not reached. Several rescue therapies have substantial inherent risks, and it is plausible that avoidance of such treatments may protect against harm (26, 27). In contrast to equivocal evidence for ECMO, a strategy that requires substantial training and resourcing, APRV may be a more accessible and cost-effective strategy (28).

In contrast to our findings, a recent RCT conducted in 52 children between the ages of 1 month to 12 years receiving mechanical ventilation for ARDS was terminated early due to harm associated with APRV (29). Generalizability to an adult population remains uncertain, and possible explanations for the differences in findings include contamination of the APRV arm and baseline imbalance favoring LTV, with lower baseline $\text{PaO}_2/\text{FiO}_2$ ratio and younger children in the APRV arm. Furthermore, compared with adults, children have higher chest wall compliance (30), and pressure-based modes such as APRV may result in greater risk of excessively large tidal volumes and risk of volutrauma although the exhaled tidal volumes were similar in the two arms of the recent pediatric RCT.

Our study has several limitations. The TSA suggests that a false positive primary outcome cannot be excluded and that larger, multicenter studies are required to substantiate our findings. All included studies were conducted in single centers, and the generalizability of the findings is uncertain, although there was no heterogeneity in the primary outcome measure.

There is a risk of study-level performance bias as blinding of the study intervention was not possible. Nonetheless, the use of mortality as an objective primary outcome measure reduces the risk of outcome assessment bias. Although APRV is conventionally defined with an end-expiratory pressure of zero, the use of positive end-expiratory pressure remains controversial. Indeed, a substantial number of the included studies used an end-expiratory pressure of greater than zero, and further exploration of the optimal delivery of APRV is required.

CONCLUSIONS

In adult patients requiring mechanical ventilation for acute hypoxic respiratory failure, APRV is associated with a mortality benefit and improved oxygenation when compared with conventional ventilation strategies. Given the limited number of patients enrolled in the available studies, larger multicenter studies are required to validate these findings.

REFERENCES

1. SRLF Trial Group: Hypoxemia in the ICU: Prevalence, treatment, and outcome. *Ann Intensive Care* 2018; 8:82
2. Brower RG, Matthay MA, Morris A, et al; The Acute Respiratory Distress Syndrome Network: Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. *N Engl J Med* 2000; 342:1301–1308
3. Cavalcanti AB, Suzumura ÉA, Laranjeira LN, et al; Writing Group for the Alveolar Recruitment for Acute Respiratory Distress Syndrome Trial (ART) Investigators: Effect of lung recruitment and titrated positive end-expiratory pressure (PEEP) vs low PEEP on mortality in patients with acute respiratory distress syndrome: A randomized clinical trial. *JAMA* 2017; 318:1335–1345
4. Brower RG, Lanken PN, MacIntyre N, et al; National Heart, Lung, and Blood Institute ARDS Clinical Trials Network: Higher versus lower positive end-expiratory pressures in patients with the acute respiratory distress syndrome. *N Engl J Med* 2004; 351:327–336
5. Mercat A, Richard JC, Vielle B, et al; Expiratory Pressure (Express) Study Group: Positive end-expiratory pressure setting in adults with acute lung injury and acute respiratory distress syndrome: A randomized controlled trial. *JAMA* 2008; 299:646–655
6. Ferguson ND, Cook DJ, Guyatt GH, et al; OSCILLATE Trial Investigators; Canadian Critical Care Trials Group: High-frequency oscillation in early acute respiratory distress syndrome. *N Engl J Med* 2013; 368:795–805
7. Habashi NM: Other approaches to open-lung ventilation: Airway pressure release ventilation. *Crit Care Med* 2005; 33:S228–S240
8. Modrykamien A, Chatburn RL, Ashton RW: Airway pressure release ventilation: An alternative mode of mechanical ventilation in acute respiratory distress syndrome. *Cleve Clin J Med* 2011; 78:101–110
9. Wrigge H, Zinserling J, Neumann P, et al: Spontaneous breathing with airway pressure release ventilation favors ventilation in dependent lung regions and counters cyclic alveolar collapse in oleic-acid-induced lung injury: A randomized controlled computed tomography trial. *Crit Care* 2005; 9:R780–R789
10. Lim J, Litton E, Robinson H, et al: Characteristics and outcomes of patients treated with airway pressure release ventilation for acute respiratory distress syndrome: A retrospective observational study. *J Crit Care* 2016; 34:154–159
11. Zhou Y, Jin X, Lv Y, et al: Early application of airway pressure release ventilation may reduce the duration of mechanical ventilation in acute respiratory distress syndrome. *Intensive Care Med* 2017; 43:1648–1659
12. Hirshberg EL, Lanspa MJ, Peterson J, et al: Randomized feasibility trial of a low tidal volume-airway pressure release ventilation

- protocol compared with traditional airway pressure release ventilation and volume control ventilation protocols. *Crit Care Med* 2018; 46:1943–1952
13. Daoud EG, Farag HL, Chatburn RL: Airway pressure release ventilation: What do we know? *Respir Care* 2012; 57:282–292
 14. Myers TR, MacIntyre NR: Respiratory controversies in the critical care setting. Does airway pressure release ventilation offer important new advantages in mechanical ventilator support? *Respir Care* 2007; 52:452–458; discussion 458–460
 15. Moher D, Liberati A, Tetzlaff J, et al; PRISMA Group: Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med* 2009; 6:e1000097
 16. Song S, Tian H, Yang X, et al: [The clinical effect of airway pressure release ventilation for acute lung injury/acute respiratory distress syndrome]. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue* 2016; 28:15–21
 17. Patel S, Sandhu R, Miller K, et al: Evaluation of airway pressure release ventilation compared to low tidal volume ventilation in ALI: Prospective Randomized Pilot Study: 425. *Crit Care Med* 2004; 32:A117
 18. Higgins JPT, Altman DG, Gøtzsche PC, et al: The cochrane collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011; 343:d5928
 19. Higgins J, Green S: *Cochrane Handbook for Systematic Reviews of Interventions*. Chichester, United Kingdom, The Cochrane Collaboration, 2011
 20. Brok J, Thorlund K, Gluud C, et al: Trial sequential analysis reveals insufficient information size and potentially false positive results in many meta-analyses. *J Clin Epidemiol* 2008; 61:763–769
 21. Li JQ, Li N, Han GJ, et al: Clinical research about airway pressure release ventilation for moderate to severe acute respiratory distress syndrome. *Eur Rev Med Pharmacol Sci* 2016; 20:2634–2641
 22. Maxwell RA, Green JM, Waldrop J, et al: A randomized prospective trial of airway pressure release ventilation and low tidal volume ventilation in adult trauma patients with acute respiratory failure. *J Trauma Acute Care Surg* 2010; 69:501–511
 23. Putensen C, Zech S, Wrigge H, et al: Long-term effects of spontaneous breathing during ventilatory support in patients with acute lung injury. *Am J Respir Crit Care Med* 2001; 164:43–49
 24. Varpula T, Valta P, Niemi R, et al: Airway pressure release ventilation as a primary ventilatory mode in acute respiratory distress syndrome. *Acta Anaesthesiol Scand* 2004; 48:722–731
 25. Ranieri VM, Rubenfeld GD, Thompson BT, et al: The ARDS Definition Task Force: Acute respiratory distress syndrome: The Berlin definition of ARDS. *JAMA* 2012; 307:2526–2533
 26. Ruan SY, Huang TM, Wu HY, et al: Inhaled nitric oxide therapy and risk of renal dysfunction: A systematic review and meta-analysis of randomized trials. *Crit Care* 2015; 19:137
 27. Vaquer S, de Haro C, Peruga P, et al: Systematic review and meta-analysis of complications and mortality of veno-venous extracorporeal membrane oxygenation for refractory acute respiratory distress syndrome. *Ann Intensive Care* 2017; 7:51
 28. Combes A, Hajage D, Capellier G, et al; EOLIA Trial Group, REVA, and ECMONet: Extracorporeal membrane oxygenation for severe acute respiratory distress syndrome. *N Engl J Med* 2018; 378:1965–1975
 29. Ganesan SL, Jayashree M, Singhi SC, et al: Airway pressure release ventilation in pediatric acute respiratory distress syndrome. A randomized controlled trial. *Am J Respir Crit Care Med* 2018; 198:1199–1207
 30. Papastamelos C, Panitch HB, England SE, et al: Developmental changes in chest wall compliance in infancy and early childhood. *J Appl Physiol (1985)* 1995; 78:179–184